

Tilburg University

The D1-triangulation of R^n for simplicial algorithms for computing solutions of nonlinear equations

Chuangyin, D.

Publication date:
1991

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Chuangyin, D. (1991). *The D1-triangulation of R^n for simplicial algorithms for computing solutions of nonlinear equations*. (Reprint Series). CentER for Economic Research.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



The D_1 -Triangulation of \mathbb{R}^n
for Simplicial Algorithms
for Computing Solutions
of Nonlinear Equations

by
Chuangyin Dang

R83
515.12

Reprinted from Mathematics of
Operations Research,
Vol. 16, No. 1, 1991

Reprint Series
no. 45

CENTER FOR ECONOMIC RESEARCH

Research Staff

Helmut Bester
Eric van Damme
Frederick van der Ploeg

Board

Helmut Bester
Eric van Damme, director
Arie Kapteyn
Frederick van der Ploeg

Scientific Council

Eduard Bomhoff	Erasmus University Rotterdam
Willem Buiters	Yale University
Jacques Drèze	Université Catholique de Louvain
Theo van de Klundert	Tilburg University
Simon Kuipers	Groningen University
Jean-Jacques Laffont	Université des Sciences Sociales de Toulouse
Merton Miller	University of Chicago
Stephen Nickell	University of Oxford
Pieter Ruys	Tilburg University
Jacques Sijben	Tilburg University

Residential Fellows

Joseph Greenberg	McGill University
Jan Magnus	Tilburg University
Emmanuel Petrakis	University of California at Los Angeles
Larry Samuelson	University of Wisconsin
Jonathan Thomas	University of Warwick

Doctoral Students

Roel Beetsma
Hans Bloemen
Chuangyin Dang
Frank de Jong
Pieter Kop Jansen

Address: Hogeschoollaan 225, P.O. Box 90153, 5000 LE Tilburg, The Netherlands
Phone : +31 13 663050
Telex : 52426 kub nl
Telefax: +31 13 663066
E-mail : "center@htikub5.bitnet"

ISSN 0924-7874

1991

Center

for

Economic Research

REPRINT

**The D_1 -Triangulation of \mathbb{R}^n
for Simplicial Algorithms
for Computing Solutions
of Nonlinear Equations**

by

Chuangyin Dang

Reprinted from Mathematics of
Operations Research,
Vol. 16, No. 1, 1991

**Reprint Series
no. 45**

THE D_1 -TRIANGULATION OF \mathbb{R}^n FOR SIMPLICIAL ALGORITHMS FOR COMPUTING SOLUTIONS OF NONLINEAR EQUATIONS*

CHUANGYIN DANG

We present a new triangulation of \mathbb{R}^n , which is called the D_1 -triangulation, for computing zero points or fixed points of nonlinear mappings. The D_1 -triangulation subdivides the unit cube and is based on very elementary pivot rules. We compare the D_1 -triangulation to several well-known triangulations of \mathbb{R}^n which triangulate the unit cube. According to several measures of efficiency the new triangulation is superior, such as the number of simplices in the unit cube, the diameter of a triangulation, the average directional density, and the surface density.

1. Introduction. There are now a number of simplicial algorithms for computing zero points or fixed points using triangulations of \mathbb{R}^n , for example, Merrill's homotopy restart method [5] and van der Laan and Talman's variable dimension simplicial restart algorithms without an extra dimension [4]. The other variable dimension algorithms have been introduced by Wright [9] and by Kojima and Yamamoto [3]. Allgower and Georg's paper [1] is an excellent survey of this field.

It has been accepted by now that the efficiency of the various simplicial homotopy and restart algorithms for solving equations is influenced in a critical manner by the triangulation employed. To evaluate and design triangulations for these algorithms, Todd, and Saigal, Solow and Wolsey established several measures in [6] and [7], such as the number of simplices in the unit cube, the diameter of a triangulation, the average directional density, and the surface density. Eaves and Yorke [2] showed that the average directional density and the surface density are equivalent.

To improve the efficiency of simplicial fixed point algorithms, we construct a new triangulation of \mathbb{R}^n and show that according to these measures it is the best of the well-known triangulations of \mathbb{R}^n , which subdivide the unit cube.

In §2 the D_1 -triangulation is introduced. We describe the pivot rules of the D_1 -triangulation in §3. The number of simplices in the unit cube, the diameter, and the surface density are calculated in §§4, 5, and 6, respectively.

2. The D_1 -triangulation of \mathbb{R}^n . Let y^0, y^1, \dots, y^k be a set of vectors in \mathbb{R}^n . If they are affinely independent, then we call their convex hull, σ , a k -simplex and write

$$\sigma = [y^0, y^1, \dots, y^k] = \text{conv}\{y^0, y^1, \dots, y^k\}.$$

A simplex τ is called a face of a simplex σ if all vertices of τ are vertices of σ . If $\dim \tau = \dim \sigma - 1$, we call τ a facet of σ . In addition, if y is the vertex of σ which is not a vertex of τ , τ is called the facet of σ opposite y .

*Received June 10, 1989; revised September 4, 1989.

AMS 1980 subject classification. Primary: 65K05. Secondary: 90C99.

IAOR 1973 subject classification. Main: Programming.

OR/MS Index 1978 subject classification. Primary: 433 Mathematics/Convexity.

Key words. Simplicial Algorithms, Triangulations, Unit Cube, Diameter, Surface Density, Average Directional Density

Let C be a convex subset of \mathbb{R}^n and let $\dim C = m$. We call G a triangulation of C if

- (1) G is a collection of m -simplices,
- (2) $C = \bigcup_{\sigma \in G} \sigma$,
- (3) for any $\sigma^1, \sigma^2 \in G$, $\sigma^1 \cap \sigma^2$ is either empty or a common face of both σ^1 and σ^2 ,

(4) each $x \in C$ has a neighborhood meeting only a finite number of simplices of G .

We denote the collection of j -simplices that are faces of simplices of G by G^j , for $j = 0, 1, \dots, m$.

For ease of notation, let $N = \{1, 2, \dots, n\}$, let $D_1^{0c} = \{y \in \mathbb{R}^n \mid \text{all components of } y \text{ are even}\}$, and for $i = 1, 2, \dots, n$, let u^i be the i th unit vector in \mathbb{R}^n .

As follows, we construct the simplices of a new triangulation of \mathbb{R}^n . We assume $n \geq 2$.

DEFINITION 2.1. Let s denote a sign vector in \mathbb{R}^n such that $s_i \in \{-1, +1\}$ for all $i \in N$. Let $0 \leq p \leq n-1$ be an integer. Let $\pi = (\pi(1), \pi(2), \dots, \pi(n))$ be a permutation of the n elements of N such that $\pi(p) < \dots < \pi(n)$ if $p \geq 1$ and $\pi(1) < \dots < \pi(n)$ if $p = 0$.

Let $y \in D_1^{0c}$. If $p = 0$, let $y^0 = y$ and

$$y^k = y + s_{\pi(k)} u^{\pi(k)}, \quad k = 1, 2, \dots, n.$$

If $p \geq 1$, let

$$y^0 = y + s,$$

$$y^k = y^{k-1} - s_{\pi(k)} u^{\pi(k)}, \quad k = 1, 2, \dots, p-1, \quad \text{and}$$

$$y^k = y + s_{\pi(k)} u^{\pi(k)}, \quad k = p, \dots, n.$$

LEMMA 2.1. Let y^0, y^1, \dots, y^n be obtained from Definition 2.1. Then y^0, y^1, \dots, y^n are affinely independent.

PROOF. If $p = 0$, then let

$$z^1 = y^1 - y^0 = s_{\pi(1)} u^{\pi(1)},$$

$$z^2 = y^2 - y^1 = s_{\pi(2)} u^{\pi(2)} - s_{\pi(1)} u^{\pi(1)},$$

...

$$z^n = y^n - y^{n-1} = s_{\pi(n)} u^{\pi(n)} - s_{\pi(n-1)} u^{\pi(n-1)}.$$

Obviously, z^1, \dots, z^n are linearly independent.

If $p \geq 1$, then let

$$z^k = y^k - y^{k-1} = -s_{\pi(k)} u^{\pi(k)}, \quad k = 1, 2, \dots, p-1,$$

$$z^p = y^p - y^{p-1} = - \sum_{k=p+1}^n s_{\pi(k)} u^{\pi(k)}, \quad \text{and}$$

$$z^k = y^k - y^{k-1} = s_{\pi(k)} u^{\pi(k)} - s_{\pi(k-1)} u^{\pi(k-1)}, \quad k = p+1, \dots, n.$$

Suppose that z^1, \dots, z^n are linearly dependent. Then there exists a $q = (q_1, \dots, q_n)^T \neq 0$ such that $q_1 z^1 + \dots + q_n z^n = 0$. If $p = n - 1$, it is necessary that $q_1 = \dots = q_{n-2} = 0$, $-q_{n-1} + q_n = 0$ and $q_n = 0$. We conclude that $q_1 = \dots = q_n = 0$. If $p < n - 1$, we must have that $q_1 = \dots = q_{p-1} = 0$, $q_{p+1} = 0$, $q_k - q_{k+1} - q_p = 0$ for $k = p + 1, \dots, n - 1$, and $q_n - q_p = 0$. Therefore, $q_p = q_n$, $q_{n-1} = 2q_n$, $q_{n-2} = 3q_n, \dots, q_{p+2} = (n - (p + 1))q_n$, and $q_{p+2} + q_p = 0$. Hence, $(n - (p + 1) + 1)q_n = 0$. Since $p < n - 1$, we have $q_1 = q_2 = \dots = q_n = 0$. Thus the hypothesis is incorrect, i.e., z^1, \dots, z^n are linearly independent. Therefore, y^0, y^1, \dots, y^n are affinely independent. The proof is completed. \square

Let y^0, y^1, \dots, y^n be obtained from Definition 2.1. Then their convex hull is an n -simplex by Lemma 2.1, which is denoted by $D_1(y, \pi, s, p)$. Let D_1 be the collection of all such simplices $D_1(y, \pi, s, p)$.

LEMMA 2.2. $\bigcup_{\sigma \in D_1} \sigma = \mathbb{R}^n$.

PROOF. Let x be an arbitrary point of \mathbb{R}^n . For each $i \in N$, let

$$y_i = \begin{cases} \lfloor x_i \rfloor & \text{if } \lfloor x_i \rfloor \text{ is even,} \\ \lfloor x_i \rfloor + 1 & \text{otherwise,} \end{cases} \quad \text{and} \quad s_i = \begin{cases} +1 & \text{if } \lfloor x_i \rfloor \text{ is even,} \\ -1, & \text{otherwise.} \end{cases}$$

We have $0 \leq \text{diag}(s_1, \dots, s_n)(x - y) \leq u$, where $u = (1, \dots, 1)^T$. Let π' be a permutation of N such that

$$0 \leq s_{\pi'(1)}(x_{\pi'(1)} - y_{\pi'(1)}) \leq \dots \leq s_{\pi'(n)}(x_{\pi'(n)} - y_{\pi'(n)}) \leq 1.$$

If $\sum_{i=1}^n s_i(x_i - y_i) \leq 1$, let

$$q'_1 = s_{\pi'(1)}(x_{\pi'(1)} - y_{\pi'(1)}), \dots, q'_n = s_{\pi'(n)}(x_{\pi'(n)} - y_{\pi'(n)}),$$

and $q'_0 = 1 - \sum_{j=1}^n q'_j$. Obviously, $q'_j \geq 0$ for all j and $\sum_{j=0}^n q'_j = 1$. Let $\pi = (1, 2, \dots, n)$, $p = 0$, $y^0 = y$, and $y^k = y + s_k u^k$ for $k = 1, 2, \dots, n$. It is easily seen that $x = \sum_{j=0}^n q'_j y^j$, where $q_0 = q'_0$ and, for $j = 1, \dots, n$, $q_j = q'_h$ with h the index for which $\pi'(h) = j$. Thus $x \in D_1(y, \pi, s, p)$.

If $\sum_{i=1}^n s_i(x_i - y_i) \geq 1$, then we show that there exists an integer $1 \leq p \leq n - 1$ such that the following system has a nonnegative solution:

$$\sum_{i=0}^{j-1} q'_i = s_{\pi'(j)}(x_{\pi'(j)} - y_{\pi'(j)}), \quad j = 1, \dots, p - 1,$$

$$\sum_{i=0}^{p-1} q'_i + q'_k = s_{\pi'(k)}(x_{\pi'(k)} - y_{\pi'(k)}), \quad k = p, \dots, n,$$

$$q'_0 + q'_1 + \dots + q'_n = 1.$$

In fact, rewriting the system, we obtain

$$\begin{aligned}
 q'_0 &= s_{\pi(1)}(x_{\pi(1)} - y_{\pi(1)}), \\
 q'_{j-1} &= s_{\pi(j)}(x_{\pi(j)} - y_{\pi(j)}) \\
 &\quad - s_{\pi(j-1)}(x_{\pi(j-1)} - y_{\pi(j-1)}), \quad j=2, \dots, p-1, \\
 q'_{p-1} &= -s_{\pi(p-1)}(x_{\pi(p-1)} - y_{\pi(p-1)}) \\
 &\quad + \left(\sum_{j=p}^n s_{\pi(j)}(x_{\pi(j)} - y_{\pi(j)}) - 1 \right) / (n-p), \\
 q'_k &= s_{\pi(k)}(x_{\pi(k)} - y_{\pi(k)}) \\
 &\quad + \left(1 - \sum_{j=p}^n s_{\pi(j)}(x_{\pi(j)} - y_{\pi(j)}) \right) / (n-p), \quad k=p, \dots, n.
 \end{aligned}$$

Let $N_0 = \{0, 1, \dots, n\}$. If $q'_{n-2} \geq 0$ for $p = n-1$, it is clear that $q'_j \geq 0$ for all $j \in N_0$; otherwise, there exists a p_0 , $1 \leq p_0 \leq n-2$, such that

$$-s_{\pi(p_0-1)}(x_{\pi(p_0-1)} - y_{\pi(p_0-1)}) + \left(\sum_{j=p_0}^n s_{\pi(j)}(x_{\pi(j)} - y_{\pi(j)}) - 1 \right) / (n-p_0) \geq 0$$

and

$$-s_{\pi(p_0)}(x_{\pi(p_0)} - y_{\pi(p_0)}) + \left(\sum_{j=p_0+1}^n s_{\pi(j)}(x_{\pi(j)} - y_{\pi(j)}) - 1 \right) / (n-p_0-1) < 0.$$

Hence,

$$\begin{aligned}
 &s_{\pi(p_0)}(x_{\pi(p_0)} - y_{\pi(p_0)}) + \left(1 - \sum_{j=p_0}^n s_{\pi(j)}(x_{\pi(j)} - y_{\pi(j)}) \right) / (n-p_0) \\
 &\geq s_{\pi(p_0)}(x_{\pi(p_0)} - y_{\pi(p_0)}) + (1 - s_{\pi(p_0)}(x_{\pi(p_0)} - y_{\pi(p_0)})) \\
 &\quad - (n-p_0-1)s_{\pi(p_0)}(x_{\pi(p_0)} - y_{\pi(p_0)}) / (n-p_0) = 0.
 \end{aligned}$$

Therefore, by taking p equal to p_0 , $q'_j \geq 0$ for all $j \in N_0$.

Let $1 \leq p \leq n-1$ be such that the system above has a nonnegative solution and let π be such that $\pi(k) = \pi'(k)$, $k = 1, 2, \dots, p-1$, and $\pi(p) < \dots < \pi(n)$.

Let

$$\begin{aligned}
 y^0 &= y + s, \\
 y^k &= y^{k-1} - s_{\pi(k)}u^{\pi(k)}, \quad k = 1, \dots, p-1, \\
 y^k &= y + s_{\pi(k)}u^{\pi(k)}, \quad k = p, \dots, n.
 \end{aligned}$$

Let q'_j be obtained from the system, for $j = 0, 1, \dots, n$. Then it is easily seen that $x = \sum_{j=0}^n q_j y^j$, where $q_0 = q'_0$ and, for $j = 1, \dots, n$, $q_j = q'_h$ with h the index for which $\pi'(h) = \pi(j)$. Thus $x \in D_1(y, \pi, s, p)$.

From these results, the lemma follows immediately. \square

LEMMA 2.3. For any σ^1 and $\sigma^2 \in D_1$, $\sigma^1 \cap \sigma^2$ is either empty or a common face of both σ^1 and σ^2 .

PROOF. Let $x \in \mathbb{R}^n$ be arbitrary. By Lemma 2.2, we may assume that $x \in \sigma$ for some

$$\sigma = [y^0, y^1, \dots, y^n] = D_1(y, \pi, s, p),$$

i.e., $x = \sum_{i=0}^n q_i y^i$, with $q_i \geq 0$ for all i and $\sum_{i=0}^n q_i = 1$. Then x lies in a face of σ whose vertices are y^j for $j \in J := \{j \in N_0 | q_j > 0\}$. We show below how each y^j , $j \in J$, can be generated from x independent of y , π , s , and p . Thus these vertices are found for any simplex of D_1 containing x .

For each $i \in N$, let

$$r_i = \begin{cases} \lfloor x_i \rfloor & \text{if } \lfloor x_i \rfloor \text{ is even,} \\ \lfloor x_i \rfloor + 1 & \text{if } \lfloor x_i \rfloor \text{ is odd,} \end{cases}$$

and

$$t_i = \begin{cases} +1 & \text{if } x_i - r_i > 0, \\ 0 & \text{if } x_i - r_i = 0, \\ -1 & \text{if } x_i - r_i < 0. \end{cases}$$

Let $w = \sum_{i=1}^n t_i(x_i - r_i)$. Further, let

$$y_i(t_j) = \begin{cases} r_i + t_j & \text{if } i = j, \\ r_i & \text{otherwise,} \end{cases}$$

for $i = 1, \dots, n$, and let $y(t_j) = (y_1(t_j), \dots, y_n(t_j))^T$. Then

$$\{y(t_1), \dots, y(t_n), r\} = \{y^j | j \in J\} \quad \text{if } w < 1, \quad \text{and}$$

$$\{y(t_1), \dots, y(t_n)\} \setminus \{r\} = \{y^j | j \in J\} \quad \text{if } w = 1.$$

Suppose that $w > 1$. Let T_1, \dots, T_g be subsets of N such that $\bigcup_{k=1}^g T_k = N$ and for each $1 \leq k \leq g$, $t_i(x_i - r_i) = t_j(x_j - r_j)$ if $i \in T_k$ and $j \in T_k$ and for any $1 \leq e < f \leq g$, $t_i(x_i - r_i) < t_j(x_j - r_j)$ if $i \in T_e$ and $j \in T_f$. Let $T_0 = \emptyset$. Let $i(k) \in T_k$ for $k = 0, \dots, g$. Since $w > 1$, there exist unique $0 \leq v < g$ and $q \geq 0$ such that

$$\begin{aligned} & t_{i(v)}(x_{i(v)} - r_{i(v)}) + (1 - |T_{v+1}| - \dots - |T_g|)q \\ & + |T_{v+1}|(t_{i(v+1)}(x_{i(v+1)} - r_{i(v+1)}) - t_{i(v)}(x_{i(v)} - r_{i(v)})) \\ & + \dots + |T_g|(t_{i(g)}(x_{i(g)} - r_{i(g)}) - t_{i(v)}(x_{i(v)} - r_{i(v)})) = 1, \end{aligned}$$

and

$$t_{i(j)}(x_{i(j)} - r_{i(j)}) - t_{i(v)}(x_{i(v)} - r_{i(v)}) - q \geq 0, \quad j = v + 1, \dots, g.$$

For $0 \leq k \leq v$, let for $i = 1, \dots, n$,

$$y_i(T_k) = \begin{cases} r_i + t_i & \text{if } i \notin T_0 \cup T_1 \cup \dots \cup T_k, \\ r_i & \text{otherwise,} \end{cases}$$

and let $y(T_k) = (y_1(T_k), \dots, y_n(T_k))^T$. For $v + 1 \leq k \leq g$, let for each $j \in T_k$,

$$\hat{y}_i(j) = \begin{cases} r_i + t_j & \text{if } i = j, \\ r_i & \text{otherwise,} \end{cases}$$

and for all i , let $\hat{y}(j) = (\hat{y}_1(j), \dots, \hat{y}_n(j))^T$. Let

$$\bar{g} = \begin{cases} g - 1 & \text{if } t_{i(j)}(x_{i(j)} - r_{i(j)}) - t_{i(v)}(x_{i(v)} - r_{i(v)}) - q = 0 \text{ for } j = g, \\ g & \text{otherwise.} \end{cases}$$

If $q = 0$, then

$$\{y(T_k) | 0 \leq k < v\} \cup \left(\bigcup_{k=v+1}^{\bar{g}} \{\hat{y}(j) | j \in T_k\} \right) = \{y^j | j \in J\},$$

and if $q > 0$, then

$$\{y(T_k) | 0 \leq k \leq v\} \cup \left(\bigcup_{k=v+1}^{\bar{g}} \{\hat{y}(j) | j \in T_k\} \right) = \{y^j | j \in J\}.$$

From these results, we obtain the proof of the lemma. \square

THEOREM 2.4. D_1 is a triangulation of \mathbb{R}^n .

PROOF. Let $x \in \mathbb{R}^n$ be arbitrary. It is clear that x is only contained in a finite number of simplices of D_1 . Using Lemma 2.1, Lemma 2.2, and Lemma 2.3, we complete the proof of the theorem. \square

The D_1 -triangulation of \mathbb{R}^3 is illustrated in Figure 1.

3. The pivot rules of the D_1 -triangulation. Let $\sigma = [y^0, y^1, \dots, y^n] = D_1(y, \pi, s, p)$ be given. We wish to obtain the unique n -simplex

$$\bar{\sigma} = [\bar{y}^0, \bar{y}^1, \dots, \bar{y}^n] = D_1(\bar{y}, \bar{\pi}, \bar{s}, \bar{p}),$$

containing all vertices of σ except y^i . Table 1 shows how \bar{y} , $\bar{\pi}$, \bar{s} , and \bar{p} depend on y , π , s , p , and i . From this table it is easy to obtain each vertex \bar{y}^k , $k = 0, 1, \dots, n$, of $\bar{\sigma}$, and in particular its new vertex.

4. Comparison of the numbers of simplices in the unit cube. Let $I^n = \{x \in \mathbb{R}^n | 0 \leq x \leq u\}$ be the unit cube in \mathbb{R}^n .

THEOREM 4.1. The number of simplices of the D_1 -triangulation in the unit cube is equal to

$$d_n = n + n(n-1) + \dots + n(n-1) \cdots 4 \cdot 3 + 2.$$




FIGURE 1. D_1 -Triangulation of the Unit Cube in \mathbb{R}^3 .

TABLE 1

The Pivot Rules of the D_1 -Triangulation

p	i	\bar{y}	\bar{s}	$\bar{\pi}$	\bar{p}
0	0	y	s	π	$p + 1$
0	$i \geq 1$	y	$s - 2s_{\pi(i)}u^{\pi(i)}$	π	p
1	0	y	s	π	$p - 1$
$2 \leq p \leq n - 1$	0	y	$s - 2s_{\pi(1)}u^{\pi(1)}$	π	p
$2 \leq p \leq n - 1$	$1 \leq i < p - 1$	y	s	$(\pi(1), \dots, \pi(i + 1), \pi(i), \dots, \pi(n))$	p
$2 \leq p \leq n - 1$	$p - 1$	y	s	$(\pi(1), \dots, \pi(p - 2), \pi(p), \dots, \pi(j), \pi(p - 1), \pi(j + 1), \dots, \pi(n))^*$	$p - 1$
$1 \leq p < n - 1$	$i > p - 1$	y	s	$(\pi(1), \dots, \pi(p - 1), \pi(i), \pi(p), \dots, \pi(i - 1), \pi(i + 1), \dots, \pi(n))$	$p + 1$
$n - 1$	$n - 1$	$y + 2s_{\pi(n)}u^{\pi(n)}$	$s - 2s_{\pi(n)}u^{\pi(n)}$	π	p
$n - 1$	n	$y + 2s_{\pi(n-1)}u^{\pi(n-1)}$	$s - 2s_{\pi(n-1)}u^{\pi(n-1)}$	π	p

*Where j is such that $\pi(j) < \pi(p - 1) < \pi(j + 1)$.

PROOF. Let $Q = \{D_1(y, \pi, s, p) | y = 0, s = (1, 1, \dots, 1)^T\}$.

From Definition 2.1, in Q , there is only one simplex for which $p = 0$, one simplex for which $p = 1$, and $n!/(n - q + 1)!$ simplices for which $p = q$, $2 \leq q \leq n - 1$. Thus

$$\begin{aligned} |Q| &= 1 + 1 + n!/(n - 1)! + n!/(n - 2)! + \dots + n!/2! \\ &= 2 + n + n(n - 1) + \dots + n(n - 1) \cdots 4 \cdot 3. \end{aligned}$$

Since $\bigcup_{\sigma \in Q} \sigma = I^n$, the proof of the theorem follows immediately. \square

For the definitions of the K_1 -, J_1 - and H_1 -triangulations, we refer to [8].

THEOREM 4.2. *The number of simplices in I^n of Freudenthal's K_1 -triangulation, that of Tucker's J_1 -triangulation, and that of Saigal's H_1 -triangulation is $n!$.*

THEOREM 4.3. *If $n \geq 3$, then $d_n < n!$. As n goes to infinity, $d_n/n!$ converges to $e - 2$.*

PROOF. For $n = 3$, we have $d_3 < 3!$, since $d_3 = 5$ and $3! = 6$. Suppose $d_{n-1} < (n - 1)!$. Thus $nd_{n-1} < n!$. From

$$\begin{aligned} nd_{n-1} &= n(n - 1) + n(n - 1)(n - 2) + \dots + n(n - 1) \cdots 4 \cdot 3 + 2n \\ &= d_n + (n - 2), \end{aligned}$$

we obtain $d_n < n!$, since $n \geq 3$. By the induction principle, the conclusion $d_n < n!$ for $n \geq 3$ follows directly. Furthermore,

$$d_n/n! = 1/(n - 1)! + 1/(n - 2)! + \dots + 1/2! + 2/n!,$$

so $d_n/n!$ converges to $e - 2$ as n goes to infinity. \square

From these results, we obtain that the number of simplices of the D_1 -triangulation is the smallest for these triangulations.

5. The diameter of the D_1 -triangulation. Let G be a triangulation of \mathbb{R}^n such that its restriction to I^n , $G|I^n = \{\sigma \subset I^n | \sigma \in G\}$, triangulates I^n and all vertices of $G|I^n$ are vertices of I^n . Let τ and τ' be two facets of G in the boundary of I^n , ∂I^n . Let $\sigma_0, \sigma_1, \dots, \sigma_m$ be a sequence of simplices of G such that σ_i and σ_{i-1} are adjacent, for $i = 1, 2, \dots, m$. If τ is a facet of σ_0 and τ' a facet of σ_m , then we say that the sequence of $\sigma_0, \sigma_1, \dots, \sigma_m$ is a path of length $m + 1$ from τ to τ' . We define the distance between τ and τ' to be the minimum length of a path between τ and τ' . The diameter of G is the maximal distance between any two facets in the boundary. It is denoted by $\text{diam}(G)$.

THEOREM 5.1.

$$\text{diam}(K_1) = 1 + n(n - 1)/2 = O(n^2),$$

$$\text{diam}(J_1) = \text{diam}(K_1),$$

$$\text{diam}(H_1) \geq (n^3 - n + 6)/6 = O(n^3), \text{ and}$$

$$\text{diam}(D_1) = 2n - 3 = O(n).$$

PROOF. Let $\sigma = [y^0, y^1, \dots, y^n] = K_1(0, \pi)$ and $\tau = [y^0, \dots, y^{n-1}]$, where $\pi = (1, 2, \dots, n)$. Let

$$\bar{\sigma} = [\bar{y}^0, \bar{y}^1, \dots, \bar{y}^n] = K_1(0, \bar{\pi}) \quad \text{and} \quad \bar{\tau} = [\bar{y}^0, \dots, \bar{y}^{n-1}],$$

where $\bar{\pi} = (n, n-1, \dots, 1)$. Let $\sigma_1, \dots, \sigma_{m-1}$ in G/I^n be such that σ_{i-1} and σ_i are adjacent for $i = 2, \dots, m-1$, σ and σ_1 are adjacent, and also σ_{m-1} and $\bar{\sigma}$. It is easily seen that the smallest m is equal to $n(n-1)/2$. The distance between τ and $\bar{\tau}$ is obviously the greatest of all distances between two facets in ∂I^n . Therefore, $\text{diam}(K_1) = n(n-1)/2 + 1$.

Since J_1/I^n is the same as K_1/I^n , $\text{diam}(J_1) = \text{diam}(K_1)$.

Let $\sigma = [y^0, y^1, \dots, y^n] = H_1(y^{2x}, \pi)$ and $\tau = [y^1, \dots, y^n]$, where $y^{2x} = (1, 0, \dots, 0)^T$ and $\pi = (1, 2, \dots, n)$. Let

$$\bar{\sigma} = [\bar{y}^0, \bar{y}^1, \dots, \bar{y}^n] = H_1(\bar{y}^{2x}, \bar{\pi}) \quad \text{and} \quad \bar{\tau} = [\bar{y}^0, \dots, \bar{y}^{n-1}],$$

where $\bar{y}^{2x} = (1, \dots, 1)^T$ and $\bar{\pi} = (n, n-1, \dots, 1)$.

Let $\sigma_1, \dots, \sigma_{m-1}$ be a sequence such that σ_{i-1} and σ_i are adjacent for $i = 2, \dots, m-1$, σ and σ_1 are adjacent, and also σ_{m-1} and $\bar{\sigma}$. Then the smallest m is equal to $(n^3 - n + 6)/6 - 1$. Thus the distance between τ and $\bar{\tau}$ is $(n^3 - n + 6)/6$. This means $\text{diam}(H_1) \geq O(n^3)$.

Finally, let $\sigma = [y^0, y^1, \dots, y^n] = D_1(y, \pi, s, p)$ and $\tau = [y^1, y^2, \dots, y^n]$, where $y = 0$, $s = (1, \dots, 1)^T$, $p = n-1$, and $\pi = (1, 2, \dots, n)$. Let

$$\bar{\sigma} = [\bar{y}^0, \bar{y}^1, \dots, \bar{y}^n] = D_1(\bar{y}, \bar{\pi}, \bar{s}, \bar{p}) \quad \text{and} \quad \bar{\tau} = [\bar{y}^1, \dots, \bar{y}^n],$$

where $\bar{y} = 0$, $\bar{s} = (1, 1, \dots, 1)^T$, $\bar{p} = n-1$, and $\bar{\pi} = (n, n-1, \dots, 3, 1, 2)$. Let $\sigma_1, \dots, \sigma_{m-1}$ be a sequence such that σ and σ_1 , σ_{i-1} and σ_i for $i = 2, \dots, m-1$, and $\bar{\sigma}$ and σ_{m-1} are adjacent. Then the smallest m is equal to $2n-4$. The distance between τ and $\bar{\tau}$ is obviously the greatest of all distances between two facets in ∂I^n . Therefore, $\text{diam}(D_1) = 2n-3$.

From these results, the theorem follows immediately. \square

6. The average directional density and surface density. From Eaves and Yorke [2], we know that for a triangulation the average directional density and surface density are equivalent. We calculate below the surface density and obtain the average directional density from the surface density.

First we calculate the surface density of the D_1 -triangulation. Let

$$\begin{aligned} \sigma^0 &= [0, u^1, \dots, u^n], & \sigma^1 &= [u, u^1, \dots, u^n], \\ \sigma^2 &= [u, u - u^1, u^2, \dots, u^n], \dots, \sigma^{n-1} \\ &= [u, u - u^1, \dots, u - u^1 - u^2 - \dots - u^{n-2}, u^{n-1}, u^n]. \end{aligned}$$

The volume of a simplex σ is denoted by $V(\sigma)$. The surface area of a simplex σ is denoted by $SA(\sigma)$. Let

$$\begin{aligned} \tau_0^0 &= [u^1, u^2, \dots, u^n], & \tau_1^0 &= [0, u^2, \dots, u^n], \dots, \tau_{n-1}^0 = [0, u^1, \dots, u^{n-2}, u^n], \\ \tau_n^0 &= [0, u^1, \dots, u^{n-1}] \end{aligned}$$

be the facets of σ^0 . Then

$$SA(\sigma^0) = \sum_{i=0}^n V(\tau_i^0) = nV(\tau_n^0) + V(\tau_0^0).$$

Clearly,

$$V(\tau_n^0) = (1/(n-1)!) |\det[u^1, u^2, \dots, u^n]| = 1/(n-1)!,$$

and

$$V(\tau_0^0) = (1/(n-1)!) |\det[u/\sqrt{n}, u^2 - u^1, \dots, u^n - u^1]| = \sqrt{n}/(n-1)!,$$

so $SA(\sigma^0) = (n + \sqrt{n})/(n-1)!$. Since $V(\sigma^0) = 1/n!$, we obtain that

$$SA(\sigma^0)/V(\sigma^0) = n(n + \sqrt{n}).$$

For $k = 2, \dots, n-1$, let

$$\tau_j^k = [u, u - u^1, \dots, u - u^1 - \dots - u^{k-1}, u^k, \dots, u^{j-1}, u^{j+1}, \dots, u^n],$$

$$j = k, \dots, n,$$

$$\tau_0^k = [u - u^1, \dots, u - u^1 - \dots - u^{k-1}, u^k, \dots, u^n], \text{ and}$$

$$\tau_j^k = [u, u - u^1, \dots, u - u^1 - \dots - u^{j-1}, u - u^1 - \dots - u^{j+1}, \dots,$$

$$u - u^1 - \dots - u^{k-1}, u^k, \dots, u^n], \quad j = 1, 2, \dots, k-1,$$

denote the facets of σ^k . Then

$$SA(\sigma^k) = V(\tau_0^k) + \sum_{j=1}^{k-1} V(\tau_j^k) + (n-k+1)V(\tau_n^k).$$

Let

$$q_1 = q_2 = \dots = q_{n-k} = ((n-k+1)^2 - 3(n-k+1) + 3)^{-1/2} \text{ and}$$

$$q_{n-k+1} = -((n-k+1)^2 - 3(n-k+1) + 3)^{-1/2}(n-k-1).$$

Then

$$V(\tau_n^k) = (1/(n-1)!) \left| \det \begin{bmatrix} 1 & 1 & \dots & 1 & 1 & \dots & 1 & 0 \\ 0 & 1 & \dots & 1 & 1 & \dots & 1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 1 & \dots & 1 & 0 \\ 0 & 0 & \dots & 0 & 0 & \dots & 1 & q_1 \\ \vdots & \vdots & & \vdots & \vdots & \ddots & \vdots & \vdots \\ \vdots & \vdots & & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 1 & \dots & 1 & q_{n-k+1} \end{bmatrix} \right|$$

$$= ((n-k+1)^2 - 3(n-k+1) + 3)^{1/2}/(n-1)!.$$

Further

$$V(\tau_0^k) = (1/(n-1)!) \det \begin{bmatrix} 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & 1 \\ 1 & 1 & \cdots & 1 & 1 & \cdots & 1 & 0 \\ 0 & 1 & \cdots & 1 & 1 & \cdots & 1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 1 & \cdots & 1 & 0 \\ 0 & 0 & \cdots & 0 & 0 & \cdots & 1 & 0 \\ \vdots & \vdots & & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 & \cdots & 0 & 0 \end{bmatrix}$$

$$= (n-k)/(n-1)!$$

Now suppose $1 \leq j \leq k-1$. If $j < k-1$, let $q_j = 1/\sqrt{2}$, $q_{j+1} = -1/\sqrt{2}$, and $q_{j+2} = \cdots = q_n = 0$. If $j = k-1$, let

$$q_{k-1} = -(n-k)((n-k+1)^2 - (n-k+1) + 1)^{-1/2} \text{ and}$$

$$q_k = \cdots = q_n = ((n-k+1)^2 - (n-k+1) + 1)^{-1/2}.$$

Then for all $j \in \{1, \dots, k-1\}$,

$$V(\tau_j^k) = (1/(n-1)!) \det \begin{bmatrix} 1 & 1 & \cdots & 1 & 1 & \cdots & 1 & 1 & \cdots & 1 & 0 \\ 0 & 1 & \cdots & 1 & 1 & \cdots & 1 & 1 & \cdots & 1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 1 & \cdots & 1 & 1 & \cdots & 1 & 0 \\ 0 & 0 & \cdots & 0 & 1 & \cdots & 1 & 1 & \cdots & 1 & q_j \\ \vdots & \vdots & & \vdots & 1 & \ddots & \vdots & \vdots & & \vdots & \vdots \\ & & & \vdots & \ddots & & 1 & & & & \\ \cdot & \cdot & & \cdot & 0 & \cdots & 1 & 1 & \cdots & 1 & q_{k-1} \\ \cdot & \cdot & & \cdot & 0 & \cdots & 0 & 0 & \cdots & 1 & q_k \\ \vdots & \vdots & & \vdots & \vdots & & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & 1 & \cdots & 0 & q_n \end{bmatrix}$$

$$= \begin{cases} (n-k)\sqrt{2}/(n-1)! & \text{if } j \neq k-1, \\ ((n-k+1)^2 - (n-k+1) + 1)^{1/2}/(n-1)! & \text{if } j = k-1. \end{cases}$$

Thus,

$$\begin{aligned} SA(\sigma^k) &= (n-k)/(n-1)! + (n-k+1) \\ &\quad \times ((n-k+1)^2 - 3(n-k+1) + 3)^{1/2}/(n-1)! \\ &\quad + (k-2)(n-k)\sqrt{2}/(n-1)! \\ &\quad + ((n-k+1)^2 - (n-k+1) + 1)^{1/2}/(n-1)!. \end{aligned}$$

Moreover,

$$V(\sigma^k) = (1/n!) \det \begin{bmatrix} 1 & 1 & \cdots & 1 & 1 & \cdots & 1 \\ 0 & 1 & \cdots & 1 & 1 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & 1 & \cdots & 1 \\ 0 & 0 & \cdots & 0 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 1 & \cdots & 0 \end{bmatrix}$$

$$= (n - k)/n!.$$

Hence,

$$\begin{aligned} SA(\sigma^k)/V(\sigma^k) &= n(n - k + (n - k + 1)((n - k + 1)^2 - 3(n - k + 1) + 3)^{1/2} \\ &\quad + (k - 2)(n - k)\sqrt{2} + ((n - k + 1)^2 \\ &\quad - (n - k + 1) + 1)^{1/2})/(n - k). \end{aligned}$$

Let

$$\begin{aligned} \tau_0^1 &= [u^1, \dots, u^n], \quad \tau_1^1 = [u, u^2, \dots, u^n], \\ \tau_2^1 &= [u, u^1, u^3, \dots, u^n], \dots, \tau_n^1 = [u, u^1, \dots, u^{n-1}] \end{aligned}$$

be the facets of σ^1 . Then

$$SA(\sigma^1) = nV(\tau_n^1) + V(\tau_0^1).$$

Let

$$q_1 = \cdots = q_{n-1} = (n^2 - 3n + 3)^{-1/2} \quad \text{and} \quad q_n = -(n - 2)(n^2 - 3n + 3)^{-1/2}.$$

Then

$$V(\tau_n^1) = (1/(n - 1)!) \det \begin{bmatrix} 0 & 1 & \cdots & 1 & q_1 \\ 1 & 0 & \cdots & 1 & q_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & \cdots & 0 & q_{n-1} \\ 1 & 1 & \cdots & 1 & q_n \end{bmatrix}$$

$$= (n^2 - 3n + 3)^{1/2}/(n - 1)!.$$

$V(\tau_0^1) = n^{1/2}/(n - 1)!$, and $V(\sigma^1) = (n - 1)/n!$. Moreover,

$$SA(\sigma^1) = (n(n^2 - 3n + 3)^{1/2} + n^{1/2})/(n - 1)!.$$

TABLE 2

Comparison of the K_I , J_I , and D_I -Triangulations

Triangulation	Number of Simplices in a Unit Cube	Diameter of a Triangulation	Average Directional Density
$K_I(J_I)$	$n!$	$O(n^2)$	$n(2 + (n-1)\sqrt{2})g_n$
D_I	$n + n(n-1) + \cdots + n(n-1)$ $\cdots 4 \cdot 3 + 2$	$O(n)$	$SD(D_I)g_n$

Hence,

$$SA(\sigma^1)/V(\sigma^1) = n(n(n^2 - 3n + 3)^{1/2} + n^{1/2})/(n-1).$$

From the above results we obtain that the surface density of the D_I -triangulation equals

$$SD(D_I) = \max\{SA(\sigma^i)/V(\sigma^i) | i = 0, 1, \dots, n-1\}.$$

Let

$$g_n = \Gamma(n/2)/((n-1)\Gamma(1/2)\Gamma((n-1)/2)).$$

From [2] we know that the average directional density of a triangulation is g_n times its surface density. Hence, the average directional density of the D_I -triangulation is equal to

$$ADD(D_I) = SD(D_I)g_n.$$

It is well known that both the average directional density of the K_I -triangulation and the one of the J_I -triangulation are equal to $n(2 + (n-1)\sqrt{2})g_n$. It is obvious that we have that $ADD(D_I) < ADD(K_I) = ADD(J_I)$, and that $ADD(D_I)/ADD(K_I)$ converges to 1 as n goes to infinity. Thus, the average directional density of the D_I -triangulation is smaller than the one of the K_I - or the J_I -triangulation. Table 2 summarizes the results above.

Acknowledgement. The author would like to thank Dolf Talman for his remarks on an earlier version of this paper, and Gerard van der Laan, He Xuchu and Chen Kaizhou for their encouragement.

References

- [1] Allgower, E. L. and Georg, K. (1980). Simplicial and Continuation Methods for Approximating Fixed Points and Solutions to Systems of Equations. *SIAM Rev.* 22 28-85.
- [2] Eaves, B. C. and Yorke, J. A. (1984). Equivalence of Surface Density and Average Directional Density. *Math. Oper. Res.* 9 363-375.
- [3] Kojima, M. and Yamamoto, Y. (1984). A Unified Approach to the Implementation of Several Restart Fixed Point Algorithms and a New Variable Dimension Algorithm. *Math. Programming* 28 288-328.
- [4] van der Laan, G. and Talman, A. J. J. (1979). A Restart Algorithm for Computing Fixed Points without an Extra Dimension. *Math. Programming* 17 74-84.

- [5] Merrill, O. H. (1972). Applications and Extensions of an Algorithm that Computes Fixed Points of Certain Upper Semi-Continuous Point to Set Mappings, Ph.D. Dissertation, Department of Industrial Engineering, University of Michigan, Ann Arbor, MI.
- [6] Saigal, R., Solow, D. and Wolsey, L. A. (1975). A Comparative Study of Two Algorithms that Compute Fixed Points in Unbounded Regions. Presented at VIIIth International Symposium on Mathematical Programming, Stanford University, Stanford, CA.
- [7] Todd, M. J. (1976). On Triangulations for Computing Fixed Points. *Math. Programming* 10 322-346.
- [8] _____ (1976). The Computation of Fixed Points and Applications. *Lecture Notes in Economics and Math. Systems* 124, Springer-Verlag, Berlin.
- [9] Wright, A. H. (1981). The Octahedral Algorithm, a New Simplicial Fixed Point Algorithm. *Math. Programming* 21 47-69.

CENTER FOR ECONOMIC RESEARCH, TILBURG UNIVERSITY, P.O. BOX 90153, 5000 LE
TILBURG, THE NETHERLANDS

Reprint Series, CentER, Tilburg University, The Netherlands:

- No. 1 G. Marini and F. van der Ploeg, Monetary and fiscal policy in an optimising model with capital accumulation and finite lives, The Economic Journal, Vol. 98, No. 392, 1988, pp. 772 - 786.
- No. 2 F. van der Ploeg, International policy coordination in interdependent monetary economies, Journal of International Economics, Vol 25, 1988, pp. 1 - 23.
- No. 3 A.P. Barten, The history of Dutch macroeconomic modelling (1936-1986), in W. Driehuis, M.M.G. Fase and H. den Hartog (eds.), Challenges for Macroeconomic Modelling, Contributions to Economic Analysis 178, Amsterdam: North-Holland, 1988, pp. 39 - 88.
- No. 4 F. van der Ploeg, Disposable income, unemployment, inflation and state spending in a dynamic political-economic model, Public Choice, Vol. 60, 1989, pp. 211 - 239.
- No. 5 Th. ten Raa and F. van der Ploeg, A statistical approach to the problem of negatives in input-output analysis, Economic Modelling, Vol. 6, No. 1, 1989, pp. 2 - 19.
- No. 6 E. van Damme, Renegotiation-proof equilibria in repeated prisoners' dilemma, Journal of Economic Theory, Vol. 47, No. 1, 1989, pp. 206 - 217.
- No. 7 C. Mulder and F. van der Ploeg, Trade unions, investment and employment in a small open economy: a Dutch perspective, in J. Muysken and C. de Neubourg (eds.), Unemployment in Europe, London: The MacMillan Press Ltd, 1989, pp. 200 - 229.
- No. 8 Th. van de Klundert and F. van der Ploeg, Wage rigidity and capital mobility in an optimizing model of a small open economy, De Economist 137, nr. 1, 1989, pp. 47 - 75.
- No. 9 G. Dhaene and A.P. Barten, When it all began: the 1936 Tinbergen model revisited, Economic Modelling, Vol. 6, No. 2, 1989, pp. 203 - 219.
- No. 10 F. van der Ploeg and A.J. de Zeeuw, Conflict over arms accumulation in market and command economies, in F. van der Ploeg and A.J. de Zeeuw (eds.), Dynamic Policy Games in Economics, Contributions to Economic Analysis 181, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1989, pp. 91 - 119.
- No. 11 J. Driffill, Macroeconomic policy games with incomplete information: some extensions, in F. van der Ploeg and A.J. de Zeeuw (eds.), Dynamic Policy Games in Economics, Contributions to Economic Analysis 181, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1989, pp. 289 - 322.
- No. 12 F. van der Ploeg, Towards monetary integration in Europe, in P. De Grauwe e.a., De Europese Monetair Integratie: vier visies, Wetenschappelijke Raad voor het Regeringsbeleid V 66, 's-Gravenhage: SDU uitgeverij, 1989, pp. 81 - 106.

- No. 13 R.J.M. Alessie and A. Kapteyn, Consumption, savings and demography, in A. Wenig, K.F. Zimmermann (eds.), Demographic Change and Economic Development, Berlin/Heidelberg: Springer-Verlag, 1989, pp. 272 - 305.
- No. 14 A. Hoque, J.R. Magnus and B. Pesaran, The exact multi-period mean-square forecast error for the first-order autoregressive model, Journal of Econometrics, Vol. 39, No. 3, 1988, pp. 327 - 346.
- No. 15 R. Alessie, A. Kapteyn and B. Melenberg, The effects of liquidity constraints on consumption: estimation from household panel data, European Economic Review 33, No. 2/3, 1989, pp. 547 - 555.
- No. 16 A. Holly and J.R. Magnus, A note on instrumental variables and maximum likelihood estimation procedures, Annales d'Économie et de Statistique, No. 10, April-June, 1988, pp. 121 - 138.
- No. 17 P. ten Hacken, A. Kapteyn and I. Woittiez, Unemployment benefits and the labor market, a micro/macro approach, in B.A. Gustafsson and N. Anders Klevmarken (eds.), The Political Economy of Social Security, Contributions to Economic Analysis 179, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1989, pp. 143 - 164.
- No. 18 T. Wansbeek and A. Kapteyn, Estimation of the error-components model with incomplete panels, Journal of Econometrics, Vol. 41, No. 3, 1989, pp. 341 - 361.
- No. 19 A. Kapteyn, P. Kooreman and R. Willemse, Some methodological issues in the implementation of subjective poverty definitions, The Journal of Human Resources, Vol. 23, No. 2, 1988, pp. 222 - 242.
- No. 20 Th. van de Klundert and F. van der Ploeg, Fiscal policy and finite lives in interdependent economies with real and nominal wage rigidity, Oxford Economic Papers, Vol. 41, No. 3, 1989, pp. 459 - 489.
- No. 21 J.R. Magnus and B. Pesaran, The exact multi-period mean-square forecast error for the first-order autoregressive model with an intercept, Journal of Econometrics, Vol. 42, No. 2, 1989, pp. 157 - 179.
- No. 22 F. van der Ploeg, Two essays on political economy: (i) The political economy of overvaluation, The Economic Journal, Vol. 99, No. 397, 1989, pp. 850 - 855; (ii) Election outcomes and the stockmarket, European Journal of Political Economy, Vol. 5, No. 1, 1989, pp. 21 - 30.
- No. 23 J.R. Magnus and A.D. Woodland, On the maximum likelihood estimation of multivariate regression models containing serially correlated error components, International Economic Review, Vol. 29, No. 4, 1988, pp. 707 - 725.
- No. 24 A.J.J. Talman and Y. Yamamoto, A simplicial algorithm for stationary point problems on polytopes, Mathematics of Operations Research, Vol. 14, No. 3, 1989, pp. 383 - 399.

- No. 25 E. van Damme, Stable equilibria and forward induction, Journal of Economic Theory, Vol. 48, No. 2, 1989, pp. 476 - 496.
- No. 26 A.P. Barten and L.J. Bettendorf, Price formation of fish: An application of an inverse demand system, European Economic Review, Vol. 33, No. 8, 1989, pp. 1509 - 1525.
- No. 27 G. Noldeke and E. van Damme, Signalling in a dynamic labour market, Review of Economic Studies, Vol. 57 (1), no. 189, 1990, pp. 1 - 23
- No. 28 P. Kop Jansen and Th. ten Raa, The choice of model in the construction of input-output coefficients matrices, International Economic Review, vol. 31, no. 1, 1990, pp. 213 - 227.
- No. 29 F. van der Ploeg and A.J. de Zeeuw, Perfect equilibrium in a model of competitive arms accumulation, International Economic Review, vol. 31, no. 1, 1990, pp. 131 - 146.
- No. 30 J.R. Magnus and A.D. Woodland, Separability and Aggregation, Economica, vol. 57, no. 226, 1990, pp. 239 - 247.
- No. 31 F. van der Ploeg, International interdependence and policy coordination in economies with real and nominal wage rigidity, Greek Economic Review, vol. 10, no. 1, June 1988, pp. 1 - 48.
- No. 32 E. van Damme, Signaling and forward induction in a market entry context, Operations Research Proceedings 1989, Berlin-Heidelberg: Springer-Verlag, 1990, pp. 45 - 59.
- No. 33 A.P. Barten, Toward a levels version of the Rotterdam and related demand systems, Contributions to Operations Research and Economics, Cambridge: MIT Press, 1989, pp. 441 - 465.
- No. 34 F. van der Ploeg, International coordination of monetary policies under alternative exchange-rate regimes, Advanced Lectures in Quantitative Economics, London-Orlando: Academic Press Ltd., 1990, pp. 91 - 121.
- No. 35 Th. van de Klundert, On socioeconomic causes of 'wait unemployment', European Economic Review, vol. 34, no. 5, 1990, pp. 1011 - 1022.
- No. 36 R.J.M. Alessie, A. Kapteyn, J.B. van Lochem and T.J. Wansbeek, Individual effects in utility consistent models of demand, in J. Hartog, G. Ridder and J. Theeuwes (eds.), Panel Data and Labor Market Studies, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1990, pp. 253 - 278.
- No. 37 F. van der Ploeg, Capital accumulation, inflation and long-run conflict in international objectives, Oxford Economic Papers, vol. 42, no. 3, 1990, pp. 501 - 525.
- No. 38 Th. Nijman and F. Palm, Parameter identification in ARMA Processes in the presence of regular but incomplete sampling, Journal of Time Series Analysis, vol. 11, no. 3, 1990, pp. 239 - 248.

- No. 39 Th. van de Klundert, Wage differentials and employment in a two-sector model with a dual labour market, Metroeconomica, vol. 40, no. 3, 1989, pp. 235 - 256.
- No. 40 Th. Nijman and M.F.J. Steel, Exclusion restrictions in instrumental variables equations, Econometric Reviews, vol. 9, no. 1, 1990, pp. 37 - 55.
- No. 41 A. van Soest, I. Woittiez and A. Kapteyn, Labor supply, income taxes, and hours restrictions in the Netherlands, Journal of Human Resources, vol. 25, no. 3, 1990, pp. 517 - 558.
- No. 42 Th.C.M.J. van de Klundert and A.B.T.M. van Schaik, Unemployment persistence and loss of productive capacity: a Keynesian approach, Journal of Macroeconomics, vol. 12, no. 3, 1990, pp. 363 - 380.
- No. 43 Th. Nijman and M. Verbeek, Estimation of time-dependent parameters in linear models using cross-sections, panels, or both, Journal of Econometrics, vol. 46, no. 3, 1990, pp. 333 - 346.
- No. 44 E. van Damme, R. Selten and E. Winter, Alternating bid bargaining with a smallest money unit, Games and Economic Behavior, vol. 2, no. 2, 1990, pp. 188 - 201.
- No. 45 C. Dang, The D_1 -triangulation of \mathbb{R}^n for simplicial algorithms for computing solutions of nonlinear equations, Mathematics of Operations Research, vol. 16, no. 1, 1991, pp. 148 - 161.

PO BOX 99450 5000 LE TILBURG THE NETHERLANDS

Bibliotheek K. U. Brabant



17 000 01169710 0